



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

$$\pi 3.2^4 R = \frac{\pi[3^{12} - 1]}{2 \log_e 3}. \quad R = \frac{3^{12} - 1}{2.3.2^4 \log_e 3}.$$

Also solved by A. H. HOLMES, J. SCHEFFER, and B. F. YANNEY.

ERRATA. In last issue, page 120, line 4 from bottom, for " $\rho = \frac{\theta^2}{c^2}$," read,

$$\rho^2 = \frac{\theta^2}{c^2}.$$

PROBLEMS.

55. Proposed by GEORGE LILLEY, Ph. D., LL. D., Principal of Park School, 394 Hall Street, Portland, Oregon.

A horse is tethered by a rope, a feet long, fastened to a post in a circular fence enclosing a circular piece of ground b feet in diameter. If the horse is outside of the fence over how much ground can he feed? If he is inside the fence over how much ground can he feed? $b > a$ in each case.

56. Proposed by Prof. B. F. BURLESON, Onside Castle, New York.

Find (1) the length s of the closed curve of the cardioid; (2) its area A ; (3) if made to revolve about its axis $2a$, find the maximum longitudinal circumference C of the solid generated; (4) find the surface K of the same; (5) its volume V ; (6) the distance x_0 of the center of gravity of the solid from the origin O ; and (7) the distance g_0 of the center of gravity of the plane curve from the origin O .

MECHANICS.

Conducted by B. F. FINKEL, Springfield, Mo. All contributions to this department should be sent to him.

SOLUTIONS OF PROBLEMS.

31. Proposed by O. W. ANTHONY, M. Sc., Professor of Mathematics in New Windsor College, New Windsor, Maryland.

A perfectly elastic, but perfectly rough mass M , and radius R , rotating in a vertical plane with an angular velocity ω , is let fall from a height, a , upon a perfectly elastic but perfectly rough horizontal plane. Determine the motion of the body after striking the plane. What will be its ultimate motion?

II. Solution by G. B. M. ZERE, A. M., Ph. D., Professor of Mathematics and Applied Science, Texarkana College, Texarkana, Arkansas-Texas.

Let V be the vertical velocity of the center just before impact; u, v , the horizontal and vertical velocities of the center just after the first impact; ω , the